

**Molloy Medical Arts Pharmacy**  
**19 Baker Avenue, Suite 207 , Poughkeepsie , NY , 12601**  
**Phone#: 8454717455      Fax#: 8454736337**

**Drug NDC:**                    **00482-0062-06**  
**Drug Name:**                **PAMINE FORTE**

**What is this medicine?**

METHSCOPOLAMINE (meth skoe POL a meen) is used to treat ulcers in the stomach and intestines. It should be given in combination with other medicines.

**This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.**

**How should I use this medicine?**

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. Take this medicine on an empty stomach, at least 30 minutes before or 2 hours after food. Do not take with food. Take your doses at regular intervals. Do not take your medicine more often than directed. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.**

**Where should I keep my medicine?**

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Throw away any unused medicine after the expiration date.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

**What should I tell my health care provider before I take this medicine?**

They need to know if you have any of these conditions: difficulty passing urine; glaucoma; heart disease; intestinal problems such as ulcerative colitis; kidney disease; liver disease; myasthenia gravis; stomach or bowel obstruction; an unusual or allergic reaction to methscopolamine, other medicines, foods, dyes, or preservatives; pregnant or trying to get pregnant; breast-feeding

**What if I miss a dose?**

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

**What should I watch for while using this medicine?**

Visit your doctor or health care professional for regular checks on your progress. You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this medicine affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. Alcohol can make you more dizzy, and increase flushing and rapid heartbeats. Avoid alcoholic drinks. Your mouth may get dry. Chewing sugarless gum or sucking hard candy, and drinking plenty of water may help. Contact your doctor if the problem does not go away or is severe. This medicine may cause dry eyes and blurred vision. If you wear contact lenses you may feel some discomfort. Lubricating drops may help. See your eye doctor if the problem does not go away or is severe. Avoid extreme heat. This medicine can cause you to sweat less than normal. Your body temperature could increase to dangerous levels, which can lead to heat stroke.

**What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible: confusion; difficulty breathing; dizziness or fainting; fast or irregular heartbeat; fever; pain or difficulty passing urine; skin rash; vomiting; weakness or tiredness. Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): anxiety, nervousness; change in sex drive or performance; constipation; difficulty sleeping; headache; loss of taste; nausea

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

**What may interact with this medicine?**

Do not take this medicine with any of the following medications: medicines used to treat erectile dysfunction (ED) like sildenafil, tadalafil, and vardenafil. This medicine may also interact with the following medications: medicines for hay fever and other allergies; medicines for mental depression; medicines for mental problems and psychotic disturbances

**This list may not describe all possible interactions. Give your health care provider a list of all the medicines, herbs, non-prescription drugs, or dietary supplements you use. Also tell them if you smoke, drink alcohol, or use illegal drugs. Some items may interact with your medicine.**