

**Molloy Medical Arts Pharmacy**  
**19 Baker Avenue, Suite 207 , Poughkeepsie , NY , 12601**  
**Phone#: 8454717455 Fax#: 8454736337**

**Drug NDC:** 52652-3001-01  
**Drug Name:** QBRELIS

**What is this medicine?**

LISINAPRIL (lyse IN oh pril) is an ACE inhibitor. This medicine is used to treat high blood pressure and heart failure. It is also used to protect the heart immediately after a heart attack.

**This medicine may be used for other purposes; ask your health care provider or pharmacist if you have questions.**

**How should I use this medicine?**

Take this medicine by mouth. Follow the directions on the prescription label. Use a specially marked spoon or container to measure each dose. Ask your pharmacist if you do not have one. Household spoons are not accurate. You may take this medicine with or without food. If it upsets your stomach, take it with food. Take your medicine at regular intervals. Do not take it more often than directed. Do not stop taking except on your doctor's advice. Talk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for children as young as 6 years for selected conditions, precautions do apply.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.** NOTE: This medicine is only for you. Do not share this medicine with others.

**Where should I keep my medicine?**

Keep out of the reach of children. Store at room temperature between 20 and 25 degrees C (68 to 77 degrees F). Throw away any unused medicine after the expiration date.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

**What should I tell my health care provider before I take this medicine?**

They need to know if you have any of these conditions: diabetes; heart or blood vessel disease; kidney disease; low blood pressure; previous swelling of the tongue, face, or lips with difficulty breathing, difficulty swallowing, hoarseness, or tightening of the throat; an unusual or allergic reaction to lisinopril, other ACE inhibitors, insect venom, food, dyes, or preservatives; pregnant or trying to get pregnant; breast-feeding.

**What if I miss a dose?**

If you miss a dose, take it as soon as you can. If it is almost time for your next dose, take only that dose. Do not take double or extra doses.

**What should I watch for while using this medicine?**

Visit your doctor or health care professional for regular check ups. Check your blood pressure as directed. Ask your doctor what your blood pressure should be, and when you should contact him or her. Do not treat yourself for coughs, colds, or pain while you are using this medicine without asking your doctor or health care professional for advice. Some ingredients may increase your blood pressure. Women should inform their doctor if they wish to become pregnant or think they might be pregnant. There is a potential for serious side effects to an unborn child. Talk to your health care professional or pharmacist for more information. Check with your doctor or health care professional if you get an attack of severe diarrhea, nausea and vomiting, or if you sweat a lot. The loss of too much body fluid can make it dangerous for you to take this medicine. You may get drowsy or dizzy. Do not drive, use machinery, or do anything that needs mental alertness until you know how this drug affects you. Do not stand or sit up quickly, especially if you are an older patient. This reduces the risk of dizzy or fainting spells. Alcohol can make you more drowsy and dizzy. Avoid alcoholic drinks. Avoid salt substitutes unless you are told otherwise by your doctor or health care professional.

**What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the hands, feet, face, lips, throat, or tongue; breathing problems; signs and symptoms of kidney injury like trouble passing urine or change in the amount of urine; signs and symptoms of increased potassium like muscle weakness; chest pain; or fast, irregular heartbeat; signs and symptoms of liver injury like dark yellow or brown urine; general ill feeling or flu-like symptoms; light-colored stools; loss of appetite; nausea; right upper belly pain; unusually weak or tired; yellowing of the eyes or skin; signs and symptoms of low blood pressure like dizziness; feeling faint or lightheaded, falls; stomach pain with or without nausea and vomiting. Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): changes in taste; cough; dizziness; fever; headache; sensitivity to light.

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

**What may interact with this medicine?**

Do not take this medicine with any of the following medications: hymenoptera venom; sacubitril; valsartan. This medicines may also interact with the following medications: aliskiren; angiotensin receptor blockers, like losartan or valsartan; certain medicines for diabetes; diuretics; everolimus; gold compounds; lithium; NSAIDs, medicines for pain and inflammation, like ibuprofen or naproxen; potassium salts or supplements; salt substitutes; sirolimus; temsirolimus.

**This list may not describe all possible interactions. Give your health care provider a list of all the medicines, herbs, non-prescription drugs, or dietary supplements you use. Also tell them if you smoke, drink alcohol, or use illegal drugs. Some items may interact with your medicine.**